


"Disobedience is the true foundation of liberty. The obedient must be slaves." ~Henry David Thoreau




Effectiveness of masks to prevent SARS-CoV-2: A Randomized Controlled Trial

Description

Conclusion: The recommendation to wear surgical masks to supplement other public health measures did not reduce the SARS-CoV-2 infection rate [*i.e.*, *no statistically significant difference between groups was found at the conventional ?-level*; $p = 0.38$; (odds ratio [OR], 0.82 [CI, 0.54 to 1.23]; $p = 0.33$)]. (Content in brackets added)

Visual summary

Image not found or type unknown



- [Download figure](#)
- [Download PowerPoint](#)

Johan Skov Bundgaard, Daniel Emil Tadeusz Raaschou-Pedersen, Christian von Buchwald, et al. [Effectiveness of Adding a Mask Recommendation to Other Public Health Measures to Prevent SARS-CoV-2 Infection in Danish Mask Wearers](#): A Randomized Controlled Trial. **Annals of Internal Medicine** 0;0 [Epub ahead of print 18 November 2020].

doi:doi.org/10.7326/M20-6817

[su_spoiler title="BibTex" style="fancy" icon="folder-2?"]
@article{doi:10.7326/M20-6817,

title = {Effectiveness of Adding a Mask Recommendation to Other Public Health Measures to Prevent SARS-CoV-2 Infection in Danish Mask Wearers},
journal = {Annals of Internal Medicine},
volume = {0},
number = {0},
pages = {null},
year = {0},
doi = {10.7326/M20-6817},
note = {PMID: 33205991},

URL = {
doi.org/10.7326/M20-6817

},
eprint = {
doi.org/10.7326/M20-6817

}
[/su_spoiler]

Abstract

Background:

Observational evidence suggests that mask wearing mitigates transmission of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2). It is uncertain if this observed association arises through protection of uninfected wearers (protective effect), via reduced transmission from infected mask wearers (source control), or both.

Objective:

To assess whether recommending surgical mask use outside the home reduces wearers' risk for SARS-CoV-2 infection in a setting where masks were uncommon and not among recommended public health measures.

Design:

Randomized controlled trial (DANMASK-19 [Danish Study to Assess Face Masks for the Protection Against COVID-19 Infection]). ([ClinicalTrials.gov](https://clinicaltrials.gov/ct2/show/study/NCT04337541): NCT04337541)

Setting:

Denmark, April and May 2020.

Participants:

Adults spending more than 3 hours per day outside the home without occupational mask use.

Intervention:

Encouragement to follow social distancing measures for coronavirus disease 2019, plus either no mask recommendation or a recommendation to wear a mask when outside the home among other persons together with a supply of 50 surgical masks and instructions for proper use.

Measurements:

The primary outcome was SARS-CoV-2 infection in the mask wearer at 1 month by antibody testing, polymerase chain reaction (PCR), or hospital diagnosis. The secondary outcome was PCR positivity for other respiratory viruses.

Results:

A total of 3030 participants were randomly assigned to the recommendation to wear masks, and 2994 were assigned to control; 4862 completed the study. Infection with SARS-CoV-2 occurred in 42 participants recommended masks (1.8%) and 53 control participants (2.1%). The between-group difference was 0.3 percentage point (95% CI, -1.2 to 0.4 percentage point; $P = 0.38$) (odds ratio, 0.82 [CI, 0.54 to 1.23]; $P = 0.33$). Multiple imputation accounting for loss to follow-up yielded similar results. Although the difference observed was not statistically significant, the 95% CIs are compatible with a 46% reduction to a 23% increase in infection.

Limitation:

Inconclusive results, missing data, variable adherence, patient-reported findings on home tests, no blinding, and no assessment of whether masks could decrease disease transmission from mask wearers to others.


Conclusion:

The recommendation to wear surgical masks to supplement other public health measures did not reduce the SARS-CoV-2 infection rate among wearers by more than 50% in a community with modest infection rates, some degree of social distancing, and uncommon general mask use. The data were compatible with lesser degrees of self-protection.

Primary Funding Source:

The Salling Foundations.

Image not found or type unknown



Fulltext: www.acpjournals.org/doi/full/10.7326/M20-6817

Category

1. General

Tags

1. COVID-19
2. Epidemiology
3. Health care providers
4. Infectious diseases
5. Lungs
6. Patients
7. Pulmonary diseases
8. SARS coronavirus
9. Upper respiratory tract infections

Date Created

29. November 2020

Author

web45